

5. Given: $\text{CO}_2(\text{g}) + \text{NaCl}(\text{aq}) + \text{NH}_3(\text{aq}) + \text{H}_2\text{O}(\text{l}) \rightarrow \text{NaHCO}_3(\text{aq}) + \text{NH}_4\text{Cl}(\text{aq})$. Determine the amount of grams of sodium bicarbonate that could be produced when 1.50 liter of carbon dioxide gas at STP is reacted with 6.11 grams NaCl, 2.70 grams of ammonia and 250.0 grams of water.
6. Given: $\text{Zn}(\text{s}) + \text{NaOH}(\text{aq}) + \text{H}_2\text{O}(\text{l}) \rightarrow \text{Na}_2\text{Zn}(\text{OH})_4(\text{aq}) + \text{H}_2(\text{g})$. What volume of hydrogen gas at STP is produced when 3.17 grams of Zn is combined with 5.00 grams of NaOH with excess water.
7. Given: $\text{CuSO}_4(\text{aq}) + \text{KI}(\text{aq}) \rightarrow \text{CuI}(\text{s}) + \text{I}_2(\text{s}) + \text{K}_2\text{SO}_4(\text{aq})$. Answer the following:
- What mass of iodine is produced when 6.76 grams of copper(II) sulfate is reacted with 9.82 grams of potassium iodide?
 - If 0.331 grams of iodine is actually produced what is the percent yield.
 - What mass of excess reactant remains?
8. What mass of lead(II) iodide is formed from the reaction of 0.357 grams of lead(II) acetate and 0.104 grams of potassium iodide?
9. A 15.32 gram sample of brass (an alloy of copper and zinc) that is 77.0% Cu by mass is reacted with 720. ml of a solution of sulfuric acid that is 14.0% H_2SO_4 by mass. What mass of CuSO_4 would be produced?
- Reaction is: $\text{Cu}(\text{s}) + \text{H}_2\text{SO}_4(\text{aq}) \rightarrow \text{CuSO}_4(\text{aq}) + \text{SO}_2(\text{g}) + \text{H}_2\text{O}(\text{l})$